

Copyright © 1996 by the American Society for Biochemistry and Molecular Biology, Inc.
9650 Rockville Pike, Bethesda, MD 20814 U.S.A.

CONTENTS Arranged by Subject Categories

CARBOHYDRATES, LIPIDS, AND OTHER
NATURAL PRODUCTS

- 9875 Expression cloning and molecular characterization of HAS protein, a eukaryotic hyaluronan synthase. Naoki Itano and Koji Kimata
- 9982 The advanced glycation end product, N^ε-(Carboxymethyl)lysine, is a product of both lipid peroxidation and glycoxidation reactions. Min-Xin Fu, Jesús R. Requena, Alicia J. Jenkins, Timothy J. Lyons, John W. Baynes, and Suzanne R. Thorpe
- 10073 A stereospecific myo-inositol/D-chiro-inositol transporter in HepG2 liver cells. Identification with D-chiro-[3-³H]inositol. Richard E. Ostlund, Jr., Robert Seemayer, Surendra Gupta, Robert Kimmel, Ellen L. Ostlund, and William R. Sherman
- 10087 Conversion from farnesyl diphosphate synthase to geranylgeranyl diphosphate synthase by random chemical mutagenesis. Shin-ichi Ohnuma, Takeshi Nakazawa, Hisashi Hemmi, Anna-Maria Hallberg, Tanetoshi Koyama, Kyoza Ogura, and Tokuzo Nishino
- 10317 α 1,3-L-fucosyltransferase expression in developing human myeloid cells. Antigenic, enzymatic, and mRNA analyses. Julia L. Clarke and Winifred M. Watkins

CELL BIOLOGY AND METABOLISM

- 9871 Glucose regulates *in vivo* glucose-6-phosphatase gene expression in the liver of diabetic rats. Duna Massillon, Nir Barzilai, Wei Chen, Meizhu Hu, and Luciano Rossetti
- 9879 Nuclear orphan receptor as a repressor of glucocorticoid receptor transcriptional activity. Thorsten Trapp and Florian Holsboer
- 9883 Regulation of insulin-like growth factor-I expression in mouse preadipocyte Ob1771 cells. Yasuki Kamai, Satoshi Mikawa, Keiji Endo, Hiroshi Sakai, and Tohru Komano
- 9891 Inhibition of p38 mitogen-activated protein kinase by insulin in cultured fetal neurons. Kim A. Heidenreich and Jennifer L. Kummer
- 9895 Sphingomyelinase and ceramide suppress insulin-induced tyrosine phosphorylation of the insulin receptor substrate-1. Hannah Kanety, Rina Hemi, Moshe Z. Papa, and Avraham Karasik
- 9902 (-)- Δ^9 -Tetrahydrocannabinol antagonizes the peripheral cannabinoid receptor-mediated inhibition of adenylyl cyclase. Michael Bayewitch, Man-Hee Rhee, Tomer Avidor-Reiss, Aviva Breuer, Raphael Mechoulam, and Zvi Vogel
- 9919 Brush border phosphatidylinositol 3-kinase mediates epidermal growth factor stimulation of intestinal NaCl absorption and Na⁺/H⁺ exchange. Seema Khurana, Samir K. Nath, Susan A. Levine, Joellyn M. Bowser, Chung-Ming Tse, Michael E. Cohen, and Mark Donowitz
- 9934 Selective uptake of cytosolic, peroxisomal, and plasma membrane proteins into the yeast lysosome for degradation. Hui-Ling Chiang, Randy Schekman, and Susan Hamamoto

- 9955 Evidence for phosphorylation of CTP:phosphocholine cytidyltransferase by multiple proline-directed protein kinases. Marcus Wieprecht, Thomas Wieder, Christine Paul, Christoph C. Geilen, and Constantin E. Orfanos
- 9969 Effects of diabetes mellitus on hepatocyte nuclear factor 1 decrease albumin gene transcription. Gonzalo Barrera-Hernandez, Irene E. Wanke, and Norman C. W. Wong
- 9976 SH2 domain function is essential for the role of the Lck tyrosine kinase in T cell receptor signal transduction. David B. Straus, Andrew C. Chan, Barbara Patai, and Arthur Weiss
- 10004 Characterization of the interaction between RhoGDI and Cdc42Hs using fluorescence spectroscopy. Tyzoon K. Nomanbhoy and Richard A. Cerione
- 10017 Reduction of BiP levels decreases heterologous protein secretion in *Saccharomyces cerevisiae*. Anne Skaja Robinson, Julie A. Bockhaus, Anne C. Voegler, and K. Dane Wittrup
- 10023 Growth-associated protein-43 (GAP-43) facilitates peptide hormone secretion in mouse anterior pituitary AtT-20 cells. Chantal Gamby, Martha C. Waage, Richard G. Allen, and Lawrence Baizer
- 10054 Association between mitogen-activated protein kinase and the ζ chain of the T cell receptor (TcR) with the SH2.3 domain of p56^{lck}. Differential regulation by TcR cross-linking. Avery August and Bo Dupont
- 10079 The degradation of human endothelial cell-derived perlecan and release of bound basic fibroblast growth factor by stromelysin, collagenase, plasmin, and heparanases. John M. Whitelock, Alan D. Murdoch, Renato V. Iozzo, and P. Anne Underwood
- 10096 Adenosine deaminase inhibition prevents free radical-mediated injury in the postschemic heart. Yong Xia, Garabet Khatchikian, and Jay L. Zweier
- 10121 Phosphorylation of Vif and its role in HIV-1 replication. Xiaoyu Yang, Joao Goncalves, and Dana Gabuzda
- 10137 Substitution of PIM1 protease in mitochondria by *Escherichia coli* Lon protease. Ulrike Teichmann, Luc van Dyck, Bernard Guiard, Heinrich Fischer, Rudi Glockshuber, Walter Neupert, and Thomas Langer
- 10149 Inactivation of Ras by *Clostridium sordellii* lethal toxin-catalyzed glucosylation. Ingo Just, Jörg Selzer, Fred Hofmann, Gaynor A. Green, and Klaus Aktories
- 10169 Oxidative stress increases production of β -amyloid precursor protein and β -amyloid (A β) in mammalian lenses, and A β has toxic effects on lens epithelial cells. Peter H. Frederikse, Donita Garland, J. Samuel Zigler, Jr., and Joram Piatigorsky
- 10183 A novel Ca²⁺-binding protein, p22, is required for constitutive membrane traffic. Margarida R. Barroso, Karen K. Bernd, Natalie D. DeWitt, Andrea Chang, Ken Mills, and Elizabeth S. Szul
- 10188 Transforming growth factors β 1, β 2, and β 3 and their receptors are differentially regulated during normal and impaired wound healing. Stefan Frank, Marianne Madlener, and Sabine Werner

•Indicates a Communication

The JBC is available on World Wide Web. URL: <http://www-jbc.stanford.edu/jbc/>

Full Instructions to Authors will be found in (1996) *J. Biol. Chem.* 271, 1-4, and reprints may be obtained from the editorial office.

- 10194 **Molecular cloning and characterization of SmLIM, a developmentally regulated LIM protein preferentially expressed in aortic smooth muscle cells.** Mukesh K. Jain, Kenji P. Fujita, Chung-Ming Hsieh, Wilson O. Endege, Nicholas E. S. Sibinga, Shau-Fang Yet, Saori Kashiki, Wen-Sen Lee, Mark A. Perrella, Edgar Haber, and Mu-En Lee
- 10200 **Insulin-mediated targeting of phosphatidylinositol 3-kinase to GLUT4-containing vesicles.** Robin A. Heller-Harrison, Michelle Morin, Adilson Guilherme, and Michael P. Czech
- 10205 **Suppression of apoptosis by dominant negative mutants of cyclin-dependent protein kinases.** William Meikrantz and Robert Schlegel
- 10217 **Ras, Rap, and Rac small GTP-binding proteins are targets for *Clostridium sordellii* lethal toxin glucosylation.** Michel R. Popoff, Esteban Chaves-Olarte, Emmanuel Lemichez, Christoph von Eichel-Streiber, Monica Thelestam, Pierre Chardin, Didier Cussac, Bruno Antonny, Philippe Chavrier, Gilles Flatau, Murielle Giry, Jean de Gunzburg, and Patrice Boquet
- 10225 **Binding of granzyme B in the nucleus of target cells. Recognition of an 80-kilodalton protein.** Michael J. Pinkoski, Ulrike Winkler, Dorothy Hudig, and R. Chris Bleakley
- 10230 **Molecular cloning and characterization of a novel human diacylglycerol kinase.** Michaeline Bunting, Wen Tang, Guy A. Zimmerman, Thomas M. McIntyre, and Stephen M. Prescott
- 10237 **Molecular cloning of a novel human diacylglycerol kinase highly selective for arachidonate-containing substrates.** Wen Tang, Michaeline Bunting, Guy A. Zimmerman, Thomas M. McIntyre, and Stephen M. Prescott
- 10256 ***Drosophila* neurocalcin, a fatty acylated, Ca^{2+} -binding protein that associates with membranes and inhibits *in vitro* phosphorylation of bovine rhodopsin.** Eva Faurobert, Ching-Kang Chen, James B. Hurley, and David Heng-Pai Teng
- 10263 **Rapid identification of yeast proteins on two-dimensional gels.** Isabelle Maillet, Gilles Lagniel, Michel Perrot, Helian Boucherie, and Jean Labarre
- 10271 **Discordant expression of osteoblast markers in MC3T3-E1 cells that synthesize a high turnover matrix.** Richard J. Wenstrup, John L. Fowlkes, David P. Witte, and Jane B. Florer
- 10277 **Demonstration of a physical interaction between microsomal triglyceride transfer protein and apolipoprotein B during the assembly of ApoB-containing lipoproteins.** Xujun Wu, Mingyue Zhou, Li-Shin Huang, John Wetterau, and Henry N. Ginsberg
- 10282 **Characterization of AMP-activated protein kinase β and γ subunits. Assembly of the heterotrimeric complex *in vitro*.** Angela Woods, Peter C. F. Cheung, Fiona C. Smith, Matthew D. Davison, James Scott, Raj K. Beri, and David Carling
- 10299 **Localization of platelet-derived growth factor-stimulated phosphorylation cascade to caveolae.** Pingsheng Liu, Yunshu Ying, Young-Gyu Ko, and Richard G. W. Anderson
- 10334 **Phosphatidic acid generation through interleukin 2 (IL-2)-induced α -diacylglycerol kinase activation is an essential step in IL-2-mediated lymphocyte proliferation.** Ignacio Flores, Teresa Casaseca, Carlos Martinez-A, Hideo Kanoh, and Isabel Merida
- 10341 **MCC, a cytoplasmic protein that blocks cell cycle progression from the G_0/G_1 to S phase.** Akihiko Matsumine, Takao Sendu, Gyeong-Hun Baeg, Badal Chandra Roy, Yusuke Nakamura, Makoto Noda, Kumao Toyoshima, and Tetsu Akiyama
- 10347 **Assembly and function of a cytosolic form of NADH-specific isocitrate dehydrogenase in yeast.** Wen-Ning Zhao and Lee McAlister-Henn
- 10372 **Partial inhibition of Na^+/K^+ -ATPase by ouabain induces the Ca^{2+} -dependent expressions of early-response genes in cardiac myocytes.** Ming Peng, Liuyu Huang, Zijian Xie, Wu-Hsiung Huang, and Amir Askari
- 10379 **Regulated cleavage of sterol regulatory element binding proteins requires sequences on both sides of the endoplasmic reticulum membrane.** Xianxin Hua, Juro Sakai, Michael S. Brown, and Joseph L. Goldstein
- 10413 **Functionally active recombinant α and β chain-peptide complexes of human major histocompatibility class II molecules.** Bishwajit Nag, Subhashini Arimilli, Prabha V. Mukku, and Irina Astafieva
- 10419 **Post-translational modification of human brain type I inositol-1,4,5-trisphosphate 5-phosphatase by farnesylation.** Florence De Smedt, Alain Boom, Xavier Pesesse, Serge N. Schiffmann, and Christophe Erney

ENZYMOLGY

- 9987 **The active species of plasma membrane Ca^{2+} -ATPase are a dimer and a monomer-calmodulin complex.** Dan L. Sackett and Danuta Kosk-Kosicka
- 10010 **Computer simulation of the triosephosphate isomerase catalyzed reaction.** Johan Aqvist and Michael Fothergill

MEMBRANES AND BIOENERGETICS

- 9898 **CD39 is an ecto- $(\text{Ca}^{2+}, \text{Mg}^{2+})$ -apyrase.** Ting-Fang Wang and Guido Guidotti
- 9906 **Photolabeling of prostaglandin endoperoxide H synthase-1 with 3-trifluoro-3-(m -[^{125}I]iodophenyl)diazirine as a probe of membrane association and the cyclooxygenase active site.** James C. Otto and William L. Smith
- 9997 **Function of the duplicated IIB domain and oligomeric structure of the fructose permease of *Escherichia coli*.** Alain Charbit, Jonathan Reizer, and Milton H. Saier, Jr.
- 10029 **Sugar binding to Na^+ /glucose cotransporters is determined by the carboxyl-terminal half of the protein.** Mariana Panayotova-Heiermann, Donald D. F. Loo, Cheng-Te Kong, Julia E. Lever, and Ernest M. Wright
- 10042 **The *ntpJ* gene in the *Enterococcus hirae ntp* operon encodes a component of KtrII potassium transport system functionally independent of vacuolar Na^+ -ATPase.** Takeshi Murata, Kazuma Takase, Ichiro Yamato, Kazuei Igarashi, and Yoshimi Kakinuma
- 10060 **Glycine betaine fluxes in *Lactobacillus plantarum* during osmotic stress and hyper- and hypo-osmotic shock.** Erwin Glaesker, Wil N. Konings, and Bert Poolman
- 10103 **Interaction of nucleotides with the NAD(H)-binding domain of the proton-translocating transhydrogenase of *Rhodospirillum rubrum*.** Tania Bizouarn, Christine Diggle, Philip G. Quirk, Rachel L. Grimley, Nick P. J. Cotton, Christopher M. Thomas, and J. Baz Jackson
- 10109 **Mutation of Tyr²³⁵ in the NAD(H)-binding subunit of the proton-translocating nicotinamide nucleotide transhydrogenase of *Rhodospirillum rubrum* affects the conformational dynamics of a mobile loop and lowers the catalytic activity of the enzyme.** Christine Diggle, Philip G. Quirk, Tania Bizouarn, Rachel L. Grimley, Nick P. J. Cotton, Christopher M. Thomas, and J. Baz Jackson
- 10210 **Identification of a new outwardly rectifying Cl^- channel that belongs to a subfamily of the ClC Cl^- channels.** Hisato Sakamoto, Masanobu Kawasaki, Shinichi Uchida, Sei Sasaki, and Fumiaki Marumo
- 10309 **Chymotryptic digestion of the cytoplasmic domain of the β subunit of Na^+/K^+ -ATPase alters kinetics of occlusion of Rb^+ ions.** Alla Shainskaya and Steven J. D. Karlisch

- 10397 **Resolution of subunit interactions and cytoplasmic sub-complexes of the yeast vacuolar proton-translocating ATPase.** John J. Tomashek, Justin L. Sonnenburg, John M. Artimovich, and Daniel J. Klionsky

NUCLEIC ACIDS, PROTEIN SYNTHESIS, AND MOLECULAR GENETICS

- 9887 **Targeted disruption of the mouse *apobec-1* gene abolishes apolipoprotein B mRNA editing and eliminates apolipoprotein B48.** Ken-ichi Hirano, Stephen G. Young, Robert V. Farese, Jr., Jennifer Ng, Eric Sande, Cara Warburton, Lyn M. Powell-Braxton, and Nicholas O. Davidson
- 9911 **Synergistic activation of transcription by the mutant and wild-type minimal transcriptional activation domain of VP16.** Subir Ghosh, Charles Toth, B. Matija Peterlin, and Edward Seto
- 9942 **AP-1/Jun is required for early *Xenopus* development and mediates mesoderm induction by fibroblast growth factor but not by activin.** Zigang Dong, Ren-He Xu, Jaebong Kim, Shu-Ning Zhan, Wei-Ya Ma, Nancy H. Colburn, and Hsiang-fu Kung
- 9947 **A regulatory element in intron 1 of the cystic fibrosis transmembrane conductance regulator gene.** Annabel N. Smith, Maria Luiza Barth, Tarra L. McDowell, Danielle S. Moulin, Hugh N. Nuthall, Michael A. Hollingsworth, and Ann Harris
- 10066 ***In vitro* interaction between human immunodeficiency virus type 1 Rev protein and splicing factor ASF/SF2-associated protein, p32.** Thomas Østergaard Tange, Torben Heick Jensen, and Jorgen Kjems
- 10143 **Involvement of G protein-coupled receptor kinase 5 in homologous desensitization of the thyrotropin receptor.** Yuji Nagayama, Kunihiro Tanaka, Takeshi Hara, Hiroyuki Namba, Shunichi Yamashita, Kohtaro Taniyama, and Masami Niwa
- 10154 **Neurotrophin-3 increases intracellular calcium in a rat insulin-secreting cell line through its action on a functional TrkC receptor.** Abdelali Tazi, Stephanie Le Bras, Hassan Ould Lamghitnia, Jean Didier Vincent, Paul Czernichow, and Raphael Scharfman
- 10242 **Multiple and tissue-specific promoter control of gonadal and non-gonadal prolactin receptor gene expression.** Zhangzhi Hu, Li Zhuang, and Maria L. Dufau
- 10291 ***In vivo* assembly of the τ -complex of the DNA polymerase III holoenzyme expressed from a five-gene artificial operon. Cleavage of the τ -complex to form a mixed γ - τ -complex by the OmpT protease.** Arthur E. Pritchard, H. Garry Dallmann, and Charles S. McHenry
- 10304 **Chicken guanylate-binding protein. Conservation of GTPase activity and induction by cytokines.** Martin Schwemmle, Bernd Kaspers, Andrea Irion, Peter Staeheli, and Ursula Schultz

- 10359 **Alternate strand DNA triple helix-mediated inhibition of HIV-1 U5 long terminal repeat integration *in vitro*.** Mohammed Bouziane, Dmitry I. Cherny, Jean-François Mouscadet, and Christian Auclair

- 10385 **Positive effect of overexpressed protein-tyrosine phosphatase PTP1C on mitogen-activated signaling in 293 cells.** Longcheng Su, Zhizhuang Zhao, Patrice Bouchard, Denis Banville, Edmond H. Fischer, Edwin G. Krebs, and Shi-Hsiang Shen

- 10391 **Visualizing a specific contact in the HIV-1 Tat protein fragment and *trans*-activation responsive region RNA complex by photocross-linking.** Yiping Liu, Zhuying Wang, and Tariq M. Rana

- 10405 **Suppression of the human erythropoietin gene expression by the TR2 orphan receptor, a member of the steroid receptor superfamily.** Han-Jung Lee, Win-Jing Young, Charles C.-Y. Shih, and Chawnsang Chang

PROTEIN CHEMISTRY AND STRUCTURE

- 9928 **Posttranslational modifications in the C-terminal tail of axonemal tubulin from sea urchin sperm.** Jean Mary, Virginie Redeker, Jean-Pierre Le Caer, Jean Rossier, and Jean-Marie Schmitter
- 9962 **Antibodies that selectively inhibit leukocyte function-associated antigen 1 binding to intercellular adhesion molecule-3 recognize a unique epitope within the CD11a I domain.** Minke E. Binnerts, Yvette van Kooyk, Caroline P. Edwards, Mark Champe, Leonard Presta, Sarah C. Bodary, Carl G. Figdor, and Philip W. Berman
- 9992 **Roles of light chains in the activity and conformation of smooth muscle myosin.** Tsuyoshi Katoh and Fumi Morita
- 10035 **Assessment of the ATP binding properties of Hsp90.** Ursula Jakob, Thomas Scheibel, Suchira Bose, Jochen Reinstein, and Johannes Buchner
- 10048 **Intracellular polymerization of the serpin plasminogen activator inhibitor type 2.** Peter Mikus and Tor Ny
- 10116 **NK-lysin, a disulfide-containing effector peptide of T-lymphocytes, is reduced and inactivated by human thioredoxin reductase. Implication for a protective mechanism against NK-lysin cytotoxicity.** Mats Andersson, Arne Holmgren, and Gianni Spyrou
- 10130 **Stability of the heme-globin linkage in $\alpha\beta$ dimers and isolated chains of human hemoglobin. A study of the heme transfer reaction from the immobilized proteins to albumin.** Maurizio Gattoni, Alberto Boffi, Paolo Sarti, and Emilia Chiancone
- 10161 **Role of aromatic transmembrane residues of the δ -opioid receptor in ligand recognition.** Katia Befort, Lina Tabbara, Dominique Kling, Bernard Maigret, and Brigitte L. Kieffer
- 10175 **Functional malleability of the carboxyl-terminal tail in protein kinase A.** Anton Chestukhin, Larisa Litovchick, Dmitry Schourov, Sarah Cox, Susan S. Taylor, and Shmuel Shaltiel

GENERAL INFORMATION

This publication is available on CD-ROM. Inquire about availability through your subscription agent or contact: THE JOURNAL OF BIOLOGICAL CHEMISTRY, P.O. Box 830399, BIRMINGHAM, AL 35283-0399, U.S.A. Minireviews are reprinted in January of the succeeding year in a Minireview Compendium. Compendia for 1988 through 1995 are available through the ASBMB office: ASBMB Office, 9650 Rockville Pike, Bethesda, MD 20814, U.S.A.

Submit all manuscripts in triplicate to

Editor, The Journal of Biological Chemistry
9650 Rockville Pike
Bethesda, MD 20814, U.S.A.

Accepted manuscripts will be published with the implicit understanding that the author(s) will pay a charge per page. Current page charges may be obtained by contacting the JBC office. Under exceptional circumstances, when no source of grant or other support exists, the author(s) may apply, at the time of submission, for a grant-in-aid to Chairman, Publications Committee, American Society for Biochemistry and Molecular Biology, Inc., 9650 Rockville Pike, Bethesda, MD 20814. All such applications must be countersigned by an appropriate institutional official stating that no funds are available for page charges.

Queries on matters of general editorial policy, requests for reprints of the "Instructions to Authors," or of the "Editorial Policy and Practices," or for

permission to reproduce any part of a previously published article should be directed to the Journal Editorial Office in Bethesda, telephone 301-530-7150; fax 301-571-1824.

The Journal of Biological Chemistry publishes papers on a broad range of topics of interest to biochemists. The views expressed are those of the author(s) and not of The Journal of Biological Chemistry or the American Society for Biochemistry and Molecular Biology.

The Journal of Biological Chemistry is copyrighted by the American Society for Biochemistry and Molecular Biology, Inc. Reprographic copying beyond that permitted by Sections 107 or 108 of the U.S. Copyright Law is allowed, provided that the \$2.50 per-copy fee is paid through the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. For those organizations that have been granted photocopy license by CCC, a separate system of payment has been arranged. The fee code for users of the Transactional Reporting Service is: 0021-9258/96/\$2.50. © Printed on acid-free paper effective with Volume 254, Issue No. 1 (1979). Reproduction of any portion of an article for subsequent republication requires permission of the copyright owner. Requests should be made in writing to the American Society for Biochemistry and Molecular Biology, Inc., Attn.: Editorial Office, 9650 Rockville Pike, Bethesda, MD 20814, and should include a statement of intended use as well as explicit specifications of the material to be reproduced.

Address all correspondence and orders relative to subscriptions and back copies to: The Journal of Biological Chemistry, P.O. Box 830399, Birmingham, AL 35283-0399, U.S.A., telephone 800-633-4931. Subscriptions are entered on a calendar year basis only. Allow at least six weeks for address changes. Claims for replacement copies must be received within three months of the issue date.

- 10247 **Cloning and sequence analysis of genes coding for *Paramecium* secretory granule (trichocyst) proteins. A unique protein fold for a family of polypeptides with different primary structures.** Marie-Christine Gautier, Linda Sperling, and Luisa Madeddu
- 10329 **Angiotensin II controls p21^{ras} activity via pp60^{c-src}.** Bernhard Schieffer, William G. Paxton, Qing Chai, Mario B. Marrero, and Kenneth E. Bernstein

- 10353 **Photolabile amiloride derivatives as cation site probes of the Na,K-ATPase.** Graham C. R. Ellis-Davies, Thomas R. Kleyman, and Jack H. Kaplan

- 10365 **The ligand recognition specificity of β_3 integrins.** Kazuhisa Suehiro, Jeffrey W. Smith, and Edward F. Plow

vii Cumulative author index. Numbers 1 through 17.

AUTHOR INDEX

- Akiyama, Tetsu, 10341
 Allmon, Klaus, 10149
 Allen, Richard G., 10023
 Anderson, Richard G. W., 10299
 Andersson, Mats, 10116
 Antony, Bruno, 10217
 Aqvist, Johan, 10010
 Arimilli, Subhashini, 10413
 Artimovich, John M., 10397
 Askari, Amir, 10372
 Astafieva, Irina, 10413
 Auclair, Christian, 10359
 August, Avery, 10054
 Avila-Jimenez, Tomer, 9902
- Baeg, Gyeong-Hun, 10341
 Baizer, Lawrence, 10023
 Banville, Denis, 10385
 Barrera-Hernandez, Gonzalo, 9969
 Barroso, Margarida R., 10183
 Barth, Maria Luiza, 9947
 Barzilai, Nir, 9871
 Bayewitch, Michael, 9902
 Baynes, John W., 9982
 Befort, Katia, 10161
 Ber, Raj K., 10282
 Bertram, Philip W., 9962
 Bernd, Karen K., 10183
 Bernstein, Kenneth E., 10329
 Binnerts, Minke E., 9962
 Bizouarn, Tania, 10103, 10109
 Bleackley, R. Chris, 10225
 Bockhaus, Julie A., 10017
 Bodary, Sarah C., 9962
 Boffi, Alberto, 10130
 Boom, Alain, 10419
 Boquet, Patrice, 10217
 Bose, Suchira, 10035
 Bourhard, Patrice, 10385
 Bouchemme, Helian, 10263
 Bouziane, Mohammed, 10359
 Bowser, Joellyn M., 9919
 Breuer, Aviva, 9902
 Brown, Michael S., 10379
 Buchner, Johannes, 10035
 Bunting, Michaeline, 10230, 10237
- Carling, David, 10282
 Casasaca, Teresa, 10334
 Cerione, Richard A., 10004
 Chai, Qing, 10329
 Champe, Mark, 9962
 Chan, Andrew C., 9976
 Chang, Andrea, 10183
 Chang, Chawnshang, 10405
 Charbit, Alain, 9997
 Chardin, Pierre, 10217
 Chaves-Orlarte, Esteban, 10217
 Chavrier, Philippe, 10217
 Chen, Ching-Kang, 10256
 Chen, Wei, 9871
 Cherny, Dmitry I., 10359
 Chestukhin, Anton, 10175
 Chisung, Peter C. F., 10282
 Chiancone, Emilia, 10130
 Chiang, Hui-Ling, 9934
 Clarke, Julia L., 10317
 Cohen, Michael E., 9919
 Colburn, Nancy H., 9942
 Cotton, Nick P. J., 10103, 10109
 Cox, Sarah, 10175
 Cussac, Didier, 10217
- Czech, Michael P., 10200
 Czernichow, Paul, 10154
- Dallmann, H. Garry, 10291
 Davidson, Nicholas O., 9887
 Davison, Matthew D., 10282
 de Gunzburg, Jean, 10217
 De Smedt, Florence, 10419
 DeWitt, Natalie D., 10183
 Diggle, Christine, 10103, 10109
 Dong, Zigang, 9942
 Donowitz, Mark, 9919
 Dufau, Maria L., 10242
 Dupont, Bo, 10054
- Edwards, Caroline P., 9962
 Ellis-Davies, Graham C. R., 10353
 Endege, Wilson O., 10194
 Endo, Keiji, 9883
 Erneux, Christophe, 10419
- Farese, Robert V., Jr., 9887
 Faurobert, Eva, 10256
 Figdor, Carl G., 9962
 Fischer, Edmond H., 10385
 Fischer, Heinrich, 10137
 Flatau, Gilles, 10217
 Florer, Jane B., 10271
 Flores, Ignacio, 10334
 Fothergill, Michael, 10010
 Fowlkes, John L., 10271
 Frank, Stefan, 10188
 Frederikse, Peter H., 10169
 Fu, Min-Xin, 9982
 Fujita, Kenji P., 10194
- Gabuzda, Dana, 10121
 Gamby, Chantal, 10023
 Garland, Donita, 10169
 Gattoni, Maurizio, 10130
 Gautier, Marie-Christine, 10247
 Geilen, Christoph C., 9955
 Ghosh, Subir, 9911
 Ginsberg, Henry N., 10277
 Giry, Murielle, 10217
 Glaesker, Erwin, 10060
 Glockshuber, Rudi, 10137
 Goldstein, Joseph L., 10379
 Goncalves, Joao, 10121
 Green, Gaynor A., 10149
 Grimley, Rachel L., 10103, 10109
 Guiard, Bernard, 10137
 Guidotti, Guido, 9898
 Guilherme, Adilson, 10200
 Gupta, Surendra, 10073
- Haber, Edgar, 10194
 Hallberg, Anna-Maria, 10087
 Hamamoto, Susan, 9934
 Hara, Takeshi, 10143
 Harris, Ann, 9947
 Heidenreich, Kim A., 9891
 Heller-Harrison, Robin A., 10200
 Hemi, Rina, 9895
 Hemmi, Hisashi, 10087
 Hirano, Ken-Ichi, 9887
 Hofmann, Fred, 10149
 Hollingsworth, Michael A., 9947
 Holmgren, Arne, 10116
 Holsboer, Florian, 9879
 Hsieh, Chung-Ming, 10194
 Hu, Meizhu, 9871
- Hu, Zhangzhi, 10242
 Hua, Xianxin, 10379
 Huang, Li-Shin, 10277
 Huang, Liuyu, 10372
 Huang, Wu-Hsiung, 10372
 Hudig, Dorothy, 10225
 Hurley, James B., 10256
- Igarashi, Kazuei, 10042
 Iozzo, Renato V., 10079
 Irion, Andrea, 10304
 Itano, Naoki, 9875
- Jackson, J. Baz, 10103, 10109
 Jain, Mukesh K., 10194
 Jakob, Ursula, 10035
 Jenkins, Alicia J., 9982
 Jensen, Torben Heick, 10066
 Just, Ingo, 10149
- Kakinuma, Yoshimi, 10042
 Kamai, Yasuki, 9883
 Kanety, Hannah, 9895
 Kanoh, Hideo, 10334
 Kaplan, Jack H., 10353
 Karasik, Avraham, 9895
 Karlsh, Steven J. D., 10309
 Kashiki, Saori, 10194
 Kaspers, Bernd, 10304
 Katoh, Tsuyoshi, 9992
 Kawasaki, Masanobu, 10210
 Khatchikian, Garabet, 10096
 Khurana, Seema, 9919
 Kieffer, Brigitte L., 10161
 Kim, Jaebong, 9942
 Kimata, Koji, 9875
 Kimmel, Robert, 10073
 Kjems, Jorgen, 10066
 Kleyman, Thomas R., 10353
 Kling, Dominique, 10161
 Klionsky, Daniel J., 10397
 Ko, Young-Gyu, 10299
 Komano, Tchrur, 9883
 Kong, Cheng-Te, 10029
 Konings, Wil N., 10060
 Kosk-Kosicka, Danuta, 9987
 Koyama, Tanetoshi, 10087
 Krebs, Edwin G., 10385
 Kummer, Jennifer L., 9891
 Kung, Hsiang-fu, 9942
- Labarre, Jean, 10263
 Lagniel, Gilles, 10263
 Lamghitnia, Hassan Ould, 10154
 Langer, Thomas, 10137
 Le Bras, Stephanie, 10154
 Le Caer, Jean-Pierre, 9928
 Lee, Han-Jung, 10405
 Lee, Mu-En, 10194
 Lee, Wen-Sen, 10194
 Lemichez, Emmanuel, 10217
 Lever, Julia E., 10029
 Levine, Susan A., 9919
 Litovchick, Larisa, 10175
 Liu, Pingsheng, 10299
 Liu, Yiping, 10391
 Loo, Donald D. F., 10029
 Lyons, Timothy J., 9982
- Ma, Wei-Ya, 9942
 Madeddu, Luisa, 10247
 Madlener, Marianne, 10188
- Maigret, Bernard, 10161
 Maillet, Isabelle, 10263
 Marrero, Mario B., 10329
 Martinez-A. Carlos, 10334
 Marumo, Fumiaki, 10210
 Mary, Jean, 9928
 Massillon, Duna, 9871
 Matsumine, Akihiko, 10341
 McAlister-Henn, Lee, 10347
 McDowell, Tarra L., 9947
 McHenry, Charles S., 10291
 McIntyre, Thomas M., 10230, 10237
- Mechoulam, Raphael, 9902
 Meikrantz, William, 10205
 Merida, Isabel, 10334
 Mikawa, Satoshi, 9883
 Mikus, Peter, 10048
 Mills, Ken, 10183
 Morin, Michelle, 10200
 Morita, Fumi, 9992
 Moulin, Danielle S., 9947
 Mouscadet, Jean-François, 10359
 Mukku, Prabha V., 10413
 Murata, Takeshi, 10042
 Murdoch, Alan D., 10079
- Nag, Bishwajit, 10413
 Nagayama, Yuji, 10143
 Nakamura, Yusuke, 10341
 Nakazawa, Takeshi, 10087
 Namba, Hiroyuki, 10143
 Nath, Samir K., 9919
 Neupert, Walter, 10137
 Ng, Jennifer, 9887
 Nishino, Tokuzo, 10087
 Niwa, Masami, 10143
 Noda, Makoto, 10341
 Nomanbhoy, Tyzoan K., 10004
 Nuthall, Hugh N., 9947
 Ny, Tor, 10048
- Ogura, Kyoze, 10087
 Ohnuma, Shin-ichi, 10087
 Orfanos, Constantin E., 9955
 Ostlund, Ellen L., 10073
 Ostlund, Richard E., Jr., 10073
 Otto, James C., 9906
- Panayotova-Heiermann, Mariana, 10029
 Papa, Moshe Z., 9895
 Patai, Barbara, 9976
 Paul, Christine, 9955
 Paxton, William G., 10329
 Peng, Ming, 10372
 Perrella, Mark A., 10194
 Perrot, Michel, 10263
 Pesesse, Xavier, 10419
 Peterlin, B. Matija, 9911
 Piatigorsky, Joram, 10169
 Pinkoski, Michael J., 10225
 Plow, Edward F., 10365
 Poolman, Bert, 10060
 Popoff, Michel R., 10217
 Powell-Braxton, Lyn M., 9887
 Prescott, Stephen M., 10230, 10237
 Presta, Leonard, 9962
 Pritchard, Arthur E., 10291
- Quirk, Philip G., 10103, 10109

CD39 Is an Ecto-(Ca²⁺, Mg²⁺)-ATPase*

(Received for publication, February 15, 1996)

Ting-Fang Wang[‡] and Guido Guidotti[§]

From the Department of Molecular and Cellular Biology, Harvard University, Cambridge, Massachusetts 02138

CD39, a 70- to 100-kDa molecule expressed primarily on activated lymphoid cells, was previously identified as a surface marker of Epstein Barr virus (EBV)-transformed B cells. In this report, we show that an ecto-(Ca²⁺, Mg²⁺)-ATPase activity is present on EBV-transformed B cells, but not on B or T lymphomas. The coincidence between CD39 expression and ecto-ATPase activity on immune cells suggests that CD39 may be an ecto-ATPase. This supposition is supported by the observation that the amino acid sequence of CD39 is significantly homologous to those of several newly identified nucleotide triphosphatases. Finally, we show that CD39 indeed has ecto-ATPase activity by expression in COS-7 cells.

Ecto-(Ca²⁺, Mg²⁺)-ATPases (ecto-ATPases) or E-type ATPases (reviewed in Ref. 1) are glycoproteins that hydrolyze extracellular nucleotide tri- and/or diphosphates. These enzymes have a high specificity activity and are insensitive to inhibitors of P-type, F-type, and V-type ATPases (2-5).

Progress in the study of ecto-ATPases has been impeded by the lack of a specific inhibitor of enzymatic activity and by their low protein abundance. Moreover, most ecto-ATPases are inactivated by the detergents normally used to solubilize membrane-bound proteins. Recently, Handa and Guidotti (6) reported that a potato tuber (*Solanum tuberosum*) ATPase was similar in amino acid sequence to CD39, a human and mouse lymphoid cell antigen (7), and to several other newly identified NTPases,¹ including yeast guanosine diphosphatase (8), garden pea nucleotide triphosphatase (NTPase),² and *Toxoplasma gondii* NTPase (10, 11). CD39, a membrane glycoprotein, was originally identified as the major surface marker of EBV-transformed B lymphoblastoid cells (12). Later it was found primarily on activated immune cells (13-15). Although the function of CD39 is largely unknown, the similarity between CD39 and the NTPases suggested that CD39 might be an ecto-ATPase.

In this report, we show that EBV-transformed B lymphocytes have ecto-ATPase activity, whereas non-EBV-trans-

formed B lymphocytes do not. Ecto-ATPase activities were also found associated with other immunocompetent cells, including cytotoxic T cells (16) and natural killer (NK) cells (4), but not with resting thymocytes and lymphomas (4, 17, 18). Both in our studies and those of other investigators, there is a coincidence between expression of CD39 and ecto-ATPase activity. All these clues lead to the hypothesis that CD39 may encode an ecto-ATPase. Finally, by transfection of COS-7 cells with CD39 cDNA, we show that CD39 indeed has ecto-ATPase activity.

MATERIALS AND METHODS

Reagents—Nucleotides, N-ethylmaleimide, ouabain, activated charcoal, DEAE-dextran, chloroquine, horseradish peroxidase-conjugated anti-mouse antibody, and nitrocellulose membrane were purchased from Sigma. [γ -³²P]ATP (triethylammonium salt) was purchased from DuPont NEN Research Products. RPMI 1640, penicillin/streptomycin, DMEM, and L-glutamine were purchased from Life Technologies, Inc. Anti-CD39 monoclonal antibody AC-2 (7, 12) was purchased from AMAC, Inc. (Westbrook, ME). Chemiluminescent reagents were purchased from Kirkegaard & Perry Laboratories (Gaithersburg, MD).

Cell Cultures—EBV-transformed B-lymphoblasts from a patient with cystic fibrosis (GM7227A) were from the Coriell Institute for Medical Research (Camden, NJ) (19). Human T leukemia Jurkat cells, EBV-transformed B lymphoblast LG2 cells, murine T lymphoma BW5147 cells, and B lymphoma A20 and Daudi cells were kindly provided by Dr. Basya Ryalov (Dept. of Molecular and Cellular Biology, Harvard University). All immune cells were maintained in medium RPMI 1640. COS-7 cells were kept at 50-70% confluence in DMEM. All cultures were supplemented with 10% fetal bovine serum, 10 mM glutamine, penicillin (100 units/ml), and streptomycin (100 μ g/ml).

Assay of ATPase Activity—ATPase activity was determined by measuring the amount of [γ -³²P]P_i released from [γ -³²P]ATP after precipitation of nucleotides with activated charcoal (4, 20). Intact cells (1 \times 10⁵) or cell homogenates were added to buffer containing 20 mM HEPES-Tris (pH 7.4), 120 mM NaCl, 5 mM KCl, 1 mM EGTA, 0.3 mM ATP, and 0.8 μ Ci of [γ -³²P]ATP; the final volume was 200 μ l. Cell extracts were obtained by homogenizing cells in the buffer with a Dounce homogenizer. The assay mixture was incubated at 37 °C for 20 min, and then the reaction was stopped by the addition of 0.5 ml of cold 20% (w/v) activated charcoal in 1.0 M HCl. The assay tubes were incubated on ice for 10 min and centrifuged at 10,000 \times g for 10 min to pellet the charcoal. Aliquots (140 μ l) of the supernatant containing the released [γ -³²P]P_i were transferred to scintillation fluid, and radioactivity was measured by using a Beckman LS5801 liquid scintillation spectrophotometer. Alternatively, the ATPase activity was determined by measuring the inorganic phosphate released as described by Ames (21), except that the time for color development was 20 min at 37 °C. All assays were performed in triplicate and reported as the mean and standard deviation. The Ca²⁺- or Mg²⁺-stimulated ATPase was determined by subtracting values obtained with EGTA alone from those with 1.5 mM CaCl₂ or MgCl₂ plus chelator.

Reverse Transcription-PCR and Construction of Expression Plasmid—Total RNA was isolated from approximately 10⁷ human EBV-transformed B lymphocytes (LG2) by the acid guanidinium thiocyanate method (22). One microgram of total RNA served as template for cDNA synthesis by avian myeloblastosis virus reverse transcriptase (Life Technologies, Inc.) for 60 min at 42 °C in the presence of 25 pmol of a specific 30-base oligonucleotide as primer in a total reaction of 20 μ l (23). The reverse transcription primer was the same one as the antisense primer to be used in the subsequent PCR reaction. The whole reverse transcription mixture was then used as template in 35 cycles of amplification. The sense primer (5'-GCCAATTCTTATGGAAGATACAAAGGAGTC) with an EcoRI site at the 5' end, contains a sequence identical with nucleotides 68-87 of human CD39 (7). The antisense primer (5'-GCTGAATTCGCTATACCATATCTTTCCAGA) is complementary to nucleotides 1581-1600 of the CD39 coding sequence, except that an EcoRI site is present. After amplification, the PCR product (1.6 kilobases) was subcloned into pCI-neo (Promega).

Expression of CD39 in COS-7 Cells and ATPase Assay—COS-7 cells

* This work was supported in part by Grant HL08893 from the National Institutes of Health. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

‡ Supported in part by a Maria Moors Cabot Fellowship.

§ To whom correspondence should be addressed: Dept. of Molecular and Cellular Biology, Harvard University, 7 Divinity Ave., Cambridge, MA 02138. Tel: 617-495-2301; Fax: 617-495-8308.

¹ The abbreviations used are: NTPase, nucleotide triphosphatase; EBV, Epstein-Barr virus; NK, natural killer cells; DMEM, Dulbecco's modified Eagle's medium; PCR, polymerase chain reaction.

² H. Hsieh and R. J. Roux, Direct submission, GenBank™ accession number Z32743.




FIG. 1. Ecto-(Ca²⁺, Mg²⁺)-apyrase activity in EBV-transformed B cells. A, time course of Ca²⁺- and Mg²⁺-stimulated apyrase activities of EBV-transformed B cell, LG2. Intact cells (left panel) or homogenate (right panel) were assayed at a concentration of 5×10^5 cells/ml in the presence of 1 mM NaN₃, 0.5 mM Na₃VO₄, and 1 mM EGTA without added Ca²⁺ or Mg²⁺ (□), or with 1.5 mM Ca²⁺ (Δ), or with 1.5 mM Mg²⁺ (○). B, ecto-(Ca²⁺)-apyrase activities of EBV-transformed B cells (LG2 and GM7227A), non-EBV-transformed B cells (A20 and Daudi), and T lymphomas (Jurkat and BW5147), assayed as described in A.

were transfected with cDNA for CD39 or with the pCI-neo vector alone by the DEAE-dextran method (24). The cells were seeded for 2 days before transfection at a density of 1700 cells per cm². On the day of transfection, the cells were washed twice with DMEM. Then, DMEM containing 10% Nuserum, 0.4 mg/ml DEAE-dextran, and chloroquine (100 μM) was added. DNA was added at a concentration of 1.25 μg/ml, and the cells were incubated for 4 h at 37 °C with 5% CO₂. Afterward, the cells were washed once with DMEM, shocked for 2 min in 10% dimethyl sulfoxide in phosphate-buffered saline and washed twice with phosphate-buffered saline. The cells were incubated in DMEM containing 10% fetal bovine serum at 37 °C with 5% CO₂. After 2 to 3 days, cells were detached by adding 10 mM EDTA and then incubated at 37 °C for 20 min. The detached cells were pooled, centrifuged, washed twice with assay buffer, and assayed for calcium-dependent ecto-apyrase activity.

Immunoblots—Cells extracts (10 or 50 μg) were resolved by 9% SDS-polyacrylamide gel electrophoresis and transferred to nitrocellulose membranes for 2 h at 500 mA. Membranes were washed in rinse buffer (phosphate-buffered saline with 3% Tween 20) at room temperature for 15 min and incubated overnight at 4 °C with AC2 anti-CD39 mAb (0.2 μg/ml) in rinse buffer. The membranes were washed three times with rinse buffer at room temperature and were incubated with a 1:2000 dilution of horseradish peroxidase-conjugated anti-mouse antibody in rinse buffer for 1.5 h at room temperature. After three washes, membranes were developed with chemiluminescent reagents, and the emitted light was recorded by x-ray film.

RESULTS

Ca²⁺ and Mg²⁺-stimulated Apyrase Activity of EBV-transformed B Lymphocytes—Human EVB-transformed B lymphocytes LG2 and GM7227A were incubated in the standard reaction buffer with [γ -³²P]ATP in the presence of 1 mM NaN₃ and 0.5 mM Na₃VO₄ for 20 min. After charcoal precipitation of the nucleotides, the released [γ -³²P]P_i in the supernatant represents the apyrase activity. The hydrolysis of [γ -³²P]ATP was linear for at least 25 min, and no radioactivity was released above background levels in assays containing no cells. Fig. 1A

TABLE I
Characterization of ecto-apyrase activity

Control ecto-apyrase activity was determined using 1×10^5 LG2 cells in the standard solution as described under "Materials and Methods." Ca²⁺-stimulated apyrase activities in the presence of inhibitors were determined at the end of a 20-min incubation by subtracting values obtained with EGTA alone from those with calcium plus chelator.

Inhibitors	% control
Control	100 ± 3
Ouabain (1 mM)	86 ± 4
VO ₄ ³⁻ (0.5 mM)	77 ± 3
N ₃ ⁻ (1 mM)	75 ± 3
N-Ethylmaleimide (10 mM)	95 ± 3
F ⁻ (10 mM)	85 ± 2

(left panel) shows the time course of Ca²⁺- and Mg²⁺-stimulated apyrase activity of intact LG2 cells. In the absence of a divalent cation, almost no hydrolysis of ATP was detected, whereas both Ca²⁺ and Mg²⁺ stimulated the activity. At the end of the incubation, cells were intact (>90%) as demonstrated by exclusion of trypan blue. Disruption of LG2 cells by homogenization in a Dounce homogenizer did not increase either Ca²⁺- or Mg²⁺-apyrase activity (right panel), suggesting that the (Ca²⁺, Mg²⁺)-apyrase activity was probably associated with the external surface of the plasma membrane. The release of P_i from EBV-transformed LG2 and GM7227A cells was 349 ± 8 and 218 ± 10 nmol per million cells per 20 min, respectively. To examine whether this hydrolytic activity was due to secreted enzymes or an intracellular enzyme released through "leaky" membrane, cells were incubated for 30 min in assay buffer without ATP and pelleted. Apyrase activity in the supernatant was less than 5% of the total apyrase activity, whereas the cell pellet contained more than 90% of the total apyrase activity. These results indicated the presence of a membrane-associated ecto-apyrase in EBV-transformed B lymphocytes.

To further characterize the observed apyrase activity, assays were done with human T leukemia Jurkat cells (ecto-apyrase-deficient (18)), African Burkitt lymphoma Daudi cells, murine T lymphoma BW5147 cells, and murine B lymphoma A20 cells. The apyrase activities of these cells were not statistically elevated above the background level (Fig. 1B).

Characterization of Ecto-apyrase Activity—Table I shows that inhibitors of P-type plasma membrane ATPases (ouabain and vanadate), V-type vacuolar ATPase (N-ethylmaleimide), and F-type mitochondrial ATPase (azide), did not significantly inhibit the EBV-transformed LG2 ecto-apyrase activity. Furthermore, 10 mM fluoride (a phosphatase inhibitor) was not inhibitory. The nucleotide specificity of the LG2 (Ca²⁺, Mg²⁺)-apyrase activity is shown in Table II. The relative hydrolysis rates of the nucleotide triphosphates is about the same, while the rate with ADP is about 60% of that with ATP. No phosphate was released from AMP (substrate of 5'-nucleotidase) and *p*-nitrophenyl phosphate (substrate of alkaline phosphatase). These results suggest that the (Ca²⁺, Mg²⁺)-apyrase activity of EBV-transformed LG2 cells is typical of an E-type ATPase.

Coincidence of CD39 Expression Pattern and Ecto-apyrase Activity in Immunocompetent Cells—CD39 was originally identified as a surface antigen of EBV-transformed lymphoblastoid cells (12). CD39 is not encoded by the EBV genome, but is a host gene. Epitope-tagging and topologic analysis indicated that CD39 has a large extracellular loop between short intracellular N and C termini. The large extracellular loop was similar in amino acid sequence to yeast guanosine phosphatase, indicating that CD39 may encode an ecto-enzyme (7). Later, CD39 was found primarily on activated immune cells, but was absent from resting thymocytes, lymphocytes, and lymphomas. In both our studies and those of other investiga-

TABLE II
Substrate specificity of ecto-(Ca²⁺)-apyrase

The apyrase activity was determined by measuring the inorganic phosphate released as described by Ames (21), except that the time for color development was 20 min at 37 °C. The LG2 cells were incubated in the standard solution with 1 mM NaN₃ and 0.5 mM Na₃VO₄. The reaction was started by adding the nucleotides to a final concentration of 1 mM and incubating at 37 °C. Ca²⁺-stimulated apyrase activity was determined at the end of a 20-min incubation by subtracting values obtained with EGTA alone from those with calcium plus chelator.

Substrates	% control
ATP	100 ± 4
GTP	89 ± 3
CTP	75 ± 3
dATP	95 ± 2
ADP	58 ± 2
AMP	0 ± 1
p-Nitrophenyl phosphate	1 ± 1

tors, there is a coincidence between the expression pattern of CD39 and ecto-apyrase activity (Table III). CD39 was only found in cells with ecto-apyrase activity, suggesting that CD39 may encode an ecto-apyrase.

Demonstration of CD39 as an Ecto-apyrase—Since detergent solubilization is a major problem in purification of many ecto-apyrases and LG2 cells lost more than 90% of the ecto-apyrase activity after solubilization with C₁₂E₉ or Triton X-100, we decided to test whether CD39 encoded an ecto-apyrase by expression. Human CD39 cDNA was amplified by reverse transcription-PCR and subcloned into the mammalian expression vector pCI-neo. COS-7 cells transfected with CD39 recombinant cDNA had about 5.4-fold higher Ca²⁺-ATPase activity (309 ± 27.6 nmol/mg/h) than cells transfected with vector alone (57.6 ± 4.8 nmol/mg/h) (Fig. 2A). This activity was not inhibited by P- or V-type ATPase inhibitors, since 1 mM NaN₃ and 0.5 mM Na₃VO₄ were included in the reaction buffer. Cells extracts (50 µg) of both preparations were examined for the presence of CD39 by Western analysis (Fig. 2B). Cells transfected with recombinant CD39 cDNA had 5-fold more CD39 protein (lane 2) as compared with cells transfected with vector alone (lane 1), estimated by comparison with the amount of CD39 protein in 10 µg (lane 3) and 50 µg (lane 4) of protein from extracts of LG2 cells.

DISCUSSION

This paper reports that CD39 is an ecto-apyrase. Four lines of evidence support this conclusion. First, the amino acid sequence of CD39 is significantly homologous to several newly identified NTPases. Second, the coincidence of the expression pattern of CD39 and ecto-apyrase activities on immunocompetent cells is consistent with the hypothesis that CD39 is an ecto-apyrase. Third, ecto-apyrase activities were found on EBV-transformed B lymphocytes that express CD39 as a major surface marker (12). No ecto-apyrase activity was found on non-EBV-transformed B lymphoma cell lines. Finally, expression of CD39 cDNA in COS-7 cells increases their ecto-apyrase activity at least 5-fold.

CD39 is expressed on activated NK cells, B cells, and T cell clones, but is not expressed by resting blood T, B, or NK cells, neutrophils, or monocytes. CD39 expression in lymphoid tissue is primarily limited to mantle zone and paracortical lymphocytes, macrophages, and dendritic cells and is generally absent from germinal centers (13). Because CD39 encodes an ecto-apyrase, the restriction of its expression to activated lymphoid cells and in anatomical sites of ongoing B cell differentiation suggests that ecto-apyrase activity may play an important role in immune responses.

One possible role of an ecto-apyrase in immune cells is to protect them from potential lytic effects of extracellular ATP

TABLE III
Distribution of ecto-apyrase activity and CD39 among immunocompetent cells

Cell lines	CD39	Ecto-apyrase
EBV-transformed B cells		
LG2	+	+
GM7727A	+	+
B cell lines		
A20	—	—
Daudi	— ^a	—
Raji	— ^{a,b}	— ^{c,d}
NK cells		
Unactivated	— ^a	ND ^e
Activated	+ ^a	+ ^e
T cell clones (activated)		
Cytotoxic cells	+ ^a	+ ^e
Helper cells	+ ^a	ND ^e
T cell lines (unactivated)		
BW5147	—	—
Jurkat	—	—
MOLT3	— ^f	— ^g
CEM	— ^{b,f}	— ^g

^a Ref. 13.

^b Ref. 12.

^c Ref. 4.

^d Ref. 18.

^e Not determined.

^f Ref. 14.

^g Ref. 17.




FIG. 2. CD39 encodes an ecto-apyrase. A, Ca²⁺-stimulated apyrase activity of COS-7 cells with expression vector pCI-neo or CD39 recombinant DNA. B, immunoblot analysis of CD39 in total cell extracts of COS-7 cells transfected with vector (lane 1, 50 µg) or CD39 recombinant DNA (lane 2, 50 µg) and of LG2 cells (lane 3, 10 µg; lane 4, 50 µg) as described under "Materials and Methods."

released by their target cells (16, 25). Extracellular ATP can induce cell death in many immune cells and a few tumor cell lines. ATP⁴⁻ binds to P₂ purinergic receptors and causes the opening of a nonselective membrane pore, which has a molecular cut-off of approximately 0.9 kDa (26). By measuring ATP-induced uptake of extracellular markers of less than 0.9 kDa, P₂ receptors were found on most immune cells and some tumor cell lines. Because of the relative high dose of extracellular ATP needed for P₂ receptor activation (half-maximal effective concentration, EC₅₀, varies from 100 µM to 1 mM), Di Virgilio (25) proposed that "leakage ATP" from the cytoplasm of stressed or injured cells was the most likely source for extracellular ATP and such an event was likely to occur at the site of inflammatory or immune responses. Hydrolysis of extracellular ATP by plasma membrane ecto-(Ca²⁺, Mg²⁺)-apyrases leads to closure of the P₂ receptor pores (27). Interestingly, persistent treatment with low doses of interleukin-2 can induce CD39 expression in T cell and NK cell *in vivo* (15); in mouse splenocytes this treatment causes resistance to extracellular ATP (28–30).

Ecto-apyrase activities have been found not only in the immune system, but also in many other tissues. The broad distribution of ecto-apyrase activity suggests that it may have

essential functions. Unfortunately, the molecular structure and function of ecto-apyrases in those tissues are largely unknown. At least three hypotheses for the roles of ecto-apyrase have been suggested. One function might be hydrolysis of ATP and other nucleotides to terminate their roles as P_2 -purinergic ligands (reviewed in Ref. 31). Secondly, ecto-apyrases may have a principal role in the formation of AMP by hydrolysis of both ADP and ATP (32). Extracellular AMP can be converted by ecto-5'-nucleotidase to adenosine, a ligand of P_1 receptors (33, 34). Both ATP and adenosine modulate a variety of key cellular processes in many tissues and organs. A third possibility is the cooperation of ecto-apyrase and 5'-nucleotidase activities with a sodium-dependent adenosine cotransporter in rat liver canalicular membranes to convert extracellular ATP to intracellular adenosine leading to conservation of the purine (9). We wonder whether CD39 or similar genes may correspond to ecto-apyrase activities in nonimmune tissues. Although CD39 was also found in endothelium (13) and human placenta,³ the expression pattern of CD39 in other nonimmune tissues has not been established.

In summary, our results show the CD39 encodes an ecto- (Ca^{2+}, Mg^{2+}) -apyrase. This is the first identification of a mammalian ecto-apyrase gene. Further analysis of CD39 may help in the understanding of the structure and function of other ecto-apyrases.

Acknowledgments—We thank Masahisa Handa for the information about the sequence similarity between the potato apyrase and CD39, Dr. Yi-Ping Hsueh for her expert advice concerning immune cell culture and for reading this manuscript, and Dr. Basya Ryalov for kindly providing the immune cell lines.

REFERENCES

1. Plesner, L. (1995) *Int. Rev. Cytol.* **158**, 1–61
2. Lin, S.-H. (1989) *J. Biol. Chem.* **264**, 14403–14407
3. Lin, S.-H. (1989) in *Cell Calcium Metabolism* (Fiskum, B., ed) pp. 13–19.
4. Dombrowski, W. E., Trevillyan, J. M., Cone, J. C., Lu, Y., and Phillips, C. A. (1993) *Biochemistry* **32**, 6515–6522
5. Stout, J. G., and Kirley, T. L. (1994) *J. Biochem. Biophys. Methods* **29**, 61–75
6. Handa, M., and Guidotti, G. (1996) *Biochem. Biophys. Res. Commun.* **218**, 916–923
7. Maliszewski, C. R., Delespesse, G. J. T., Schoenborn, M. A., Armitage, R. J., Fanslow, W. C., Nakajima, T., Baker, E., Sutherland, G. R., Poindexter, K., Birks, C., Alpert, A., Friend, D., Gimpel, S. D., and Gayle, R. B., III (1994) *J. Immunol.* **153**, 3574–3583
8. Abeijon, C., Yanagisawa, K., Mandon, E. C., Hauser, A., Moremen, C. B., Hirschberg, C. B., and Robbins, P. W. (1993) *J. Cell Biol.* **122**, 307–313
9. Chen, M., Nishida, T., Gatmaitan, Z., and Arias, I. M. (1992) *J. Biol. Chem.* **267**, 9684–9688
10. Bermude, D., Peck, K. R., Afifi, M. A., Becker, C. J. M., and Joiner, K. A. (1994) *J. Biol. Chem.* **269**, 29252–29260
11. Asai, T., Miura, S., Sibley, L. D., Okabayashi, H., and Takeuchi, T. (1995) *J. Biol. Chem.* **270**, 11391–11397
12. Rowe, M., Hildreth, J. E. K., Rickinson, A. B., and Epstein, M. A. (1982) *Int. J. Cancer* **29**, 373–381
13. Kansas, G. S., Wood, G. S., and Tedder, T. F. (1991) *J. Immunol.* **146**, 2235–2244
14. Ikewaki, N., Takabe, S., and Tsuji, K. (1992) *Tissue Antigen* **39**, 174–181
15. Deunsing, S., Kirsher, H., and Atzpodien, J. (1994) *Blood* **83**, 3826–3827
16. Fillipini, A., Taffa, R. E., Agui, T., and Sitkovsky, M. V. (1990) *J. Biol. Chem.* **267**, 16081–16087
17. Barankiewicz, J., Dosch, H.-M., and Cohen, A. (1988) *J. Biol. Chem.* **263**, 7094–7098
18. Barankiewicz, J., and Cohn, A. (1990) *Ann. N. Y. Acad. Sci.* **603**, 380–393
19. Chen, H., Schulman, H., and Garder, P. (1989) *Science* **243**, 657–660
20. Reddy, N. B., Oliver, K. L., Festoff, B. W., and Engle, W. K. (1978) *Biochim. Biophys. Acta* **540**, 371–388
21. Ames, B. N. (1966) *Methods Enzymol.* **8**, 115–117
22. Chomczynski, P., and Sacchi, N. (1987) *Anal. Biochem.* **162**, 156–159
23. Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd Ed, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
24. Lopata, M. A., Cleveland, D. M., and Sollner-Webb, B. (1984) *Nucleic Acids Res.* **12**, 5707–5717
25. Di Virgilio, F. (1995) *Immunol. Today* **16**, 524–528
26. Di Virgilio, F., Bronte, V., Collavo, D., and Zanovello, P. (1989) *J. Immunol.* **143**, 1955–1960
27. Di Virgilio, F., and Steinberg, T. H. (1993) *Cell and Tissue Culture: Laboratory Procedures*, Vol. 5B, pp. 4.1–4.8, John Wiley and Sons, New York
28. Zanovello, P., Bronte, V., Pizzo, P., and Di Virgilio, F. (1990) *J. Immunol.* **145**, 1545–1550
29. Zheng, L. M., Zychlinsky, A., Liu, C.-C., Ojcius, D. M., and Young, J. D.-E. (1991) *J. Cell Biol.* **112**, 279–288
30. Pizzo, P., Murgia, M., and Zamboni, A. (1992) *J. Immunol.* **149**, 3372–3378
31. Dubyak, G. R., and El-Motassim, C. (1993) *Am. J. Physiol.* **34**, C577–C606
32. Lin, S.-H. (1985) *J. Biol. Chem.* **260**, 10976–10980
33. Zimmermann, H. (1992) *Biochem. J.* **285**, 345–365
34. Olah, M. E., and Stiles, G. L. (1995) *Annu. Rev. Pharmacol. Toxicol.* **35**, 518–606

³ L. Hillier, N. Clark, T. Dubuque, K. Elliston, M. Hawkins, M. Holman, M. Hultman, T. Kucaba, M. Le, G. Lennon, M. Marra, J. Parson, L. Rifkin, T. Rohlfing, M. Soares, F. Tan, E. Trevaskis, R. Waterston, A. Williamson, P. Wohldmann, and R. Wilson, Direct submission, GenBank™ accession numbers H13211 and H13577.